

Non-invasive brain stimulation: a new strategy to improve neurorehabilitation after stroke?

Friedhelm C Hummel, Leonardo G Cohen

Lancet Neurol 2006; 5: 708–12

Human Cortical Physiology
Section and Stroke
Neurorehabilitation Clinic,
National Institute of
Neurological Disorders and
Stroke, National Institute of
Health, Bethesda, USA
(L G Cohen MD, C Hummel MD);
and Cortical Physiology
Research Group, Department of
Neurology, Hamburg
University Medical Center,
Hamburg, Germany
(F C Hummel)

Correspondence to:
Leonardo G Cohen, Human
Cortical Physiology Section,
NINDS, NIH, Bethesda,
MD 20817, USA
cohenl@ninds.nih.gov

Background Motor impairment resulting from chronic stroke can have extensive physical, psychological, financial, and social implications despite available neurorehabilitative treatments. Recent studies in animals showed that direct epidural stimulation of the primary motor cortex surrounding a small infarct in the lesioned hemisphere ($M1_{\text{lesioned hemisphere}}$) elicits improvements in motor function.

Recent developments In human beings, proof of principle studies from different laboratories showed that non-invasive transcranial magnetic stimulation and direct current stimulation that upregulate excitability within $M1_{\text{lesioned hemisphere}}$ or downregulate excitability in the intact hemisphere ($M1_{\text{intact hemisphere}}$) results in improvement in motor function in patients with stroke. Possible mechanisms mediating these effects can include the correction of abnormally persistent interhemispheric inhibitory drive from $M1_{\text{intact hemisphere}}$ to $M1_{\text{lesioned hemisphere}}$ in the process of generation of voluntary movements by the paretic hand, a disorder correlated with the magnitude of impairment. In this paper we review these mechanistically oriented interventional approaches.

What next? These findings suggest that transcranial magnetic stimulation and transcranial direct current stimulation could develop into useful adjuvant strategies in neurorehabilitation but have to be further assessed in multicentre clinical trials.

Introduction

Recovery of motor function after stroke is typically incomplete.¹ 6 months after the episode, two-thirds of stroke survivors are unable to take part in activities of daily living with their paretic hand to the extent they were before² and only a few are able to carry out professional work.³ It remains a desirable goal to develop strategies to improve the beneficial effects of neurorehabilitative treatments.

Neuroimaging studies showed increased activity of $M1_{\text{intact hemisphere}}$ with movements of the paretic hand in patients with motor impairment.^{4,5} The role of activity in the intact hemisphere on motor control, presently under investigation, varies depending on lesion site, time from stroke, and magnitude of impairment.^{6,7} Patients with stroke experience changes in motor cortical excitability^{8–11} and an abnormally high interhemispheric inhibition from $M1_{\text{intact hemisphere}}$ to $M1_{\text{lesioned hemisphere}}$ with movements of the paretic hand that is more prominent in cases with more substantial motor impairment.¹ These findings, consistent with interhemispheric competition models of sensory and motor processing,^{12–15} raised the hypothesis that purposeful modulation of excitability in motor regions of the intact and affected hemisphere may contribute to improvements in motor function.¹⁶

Recent studies in animals showed that motor recovery after focal lesions in the hand motor representation can improve with direct epidural cortical stimulation.^{17–19} The feasibility and safety of this invasive approach in patients is under investigation;²⁰ an alternative approach is the use of non-invasive cortical stimulation.

Non-invasive brain stimulation is a powerful method to modulate human brain function.^{21–23} Transcranial

magnetic stimulation is a painless procedure that modulates cortical excitability and has contributed to the understanding of mechanisms underlying cognitive processes.^{21,24} The procedure involves a short strong electrical current that is delivered through an insulated coil of wire placed over the scalp (magnetic coil). The induced electrical currents modulate neuronal excitability at the stimulated sites. Depending on stimulation parameters, transcranial magnetic stimulation can upregulate or downregulate excitability to different extents²⁵ in the neural structures under the stimulating coil.²⁶ Transcranial direct-current stimulation is a procedure used to polarise brain regions through the non-invasive application of weak direct currents.^{23,27,28} The procedure elicits focal reversible shifts in cortical excitability depending on the polarity, strength, and duration of stimulation.²⁸ Both techniques can purposefully modulate brain function, are painless and non-invasive, and can be used in double-blind, experimental designs^{27,29–34}—although the mechanisms underlying these features may differ (panel^{22,23,27,35–38}).

Interhemispheric competition models¹⁶ suggest possible strategies to influence function in the paretic hand (figure): upregulation of excitability in $M1_{\text{lesioned hemisphere}}$ and downregulation of excitability in $M1_{\text{intact hemisphere}}$. In this paper we describe recent studies that report the effects of both transcranial magnetic stimulation and transcranial direct current stimulation on cortical excitability and motor function in the paretic hand after chronic stroke. Additional options under investigation include modulation of excitability in non-primary motor regions like the dorsal^{39,40} and ventral⁴¹ premotor cortices or the supplementary motor area.

Recent developments

Upregulation of excitability in the affected hemisphere

Two non-invasive strategies have been used to increase excitability in the affected hemisphere: anodal transcranial direct current stimulation and rapid-rate transcranial magnetic stimulation (panel^{24,26,28,35}). Anodal transcranial direct current stimulation delivered to M1_{lesioned hemisphere} was studied in patients with chronic stroke in sham-controlled double-blind crossover experimental designs.^{29–31} These studies showed improvements in performance of motor tasks that mimic activities of daily living (eg, Jebsen Taylor hand function test)⁴² with treatment but not with sham stimulation that lasted for more than 30 min after the end of the stimulation period.²⁹ Performance of less complex motor tasks, such as simple force generation and reaction time tasks, is improved in similar ways (Hummel and colleagues, unpublished data). Behavioural gains were accompanied by enhanced cortical excitability and reduced intracortical inhibition^{29,30} within M1_{lesioned hemisphere}, suggesting the involvement of glutamatergic and GABAergic neurotransmission as possible operating mechanisms.

Khedr and colleagues³² applied repetitive transcranial magnetic stimulation daily (ten 10 s trains of 3 Hz stimulation with 50 s between each train) combined with customary rehabilitative treatment for 10 days within the first 2 weeks after stroke. The researchers reported motor improvements with repetitive transcranial magnetic stimulation relative to sham lasting for at least 10 days after the end of the treatment.³² There were no complications with either interventions and the researchers proposed that repetitive stimulation sessions could elicit longer-lasting effects than single applications.

Downregulation of excitability in the intact hemisphere

An alternative approach that could theoretically improve motor function in the paretic hand is downregulation of excitability in M1_{intact hemisphere}, possibly through modulation of inappropriate interhemispheric inhibition.¹ Previous studies in healthy volunteers showed that downregulation of excitability in one motor cortex results in increased excitability in the opposite motor cortex^{43,44} and performance improvements in motor functions of the ipsilateral hand,⁴⁵ a mechanism proposed to rely on modulation of interhemispheric inhibition between both primary motor cortices.^{44,46} Two studies recently tested this hypothesis applying low-frequency repetitive transcranial magnetic stimulation to M1_{intact hemisphere} after chronic stroke in double-blind sham-controlled experimental designs.^{33,34} Mansur and colleagues³³ used a crossover design in their study, whereas Takeuchi and colleagues³⁴ applied real repetitive transcranial magnetic stimulation in one group of patients and sham stimulation in another group. Both studies reported improvements in motor functions in the paretic hand, characterised by improved reaction times and pegboard

Panel: Non-invasive cortical stimulation

Transcranial magnetic stimulation (TMS) and transcranial direct-current stimulation (tDCS) modulate cortical activity and influence motor, sensory, and cognitive functions

TMS

Delivered to the brain by passing a strong brief electrical current through an insulated wired coil placed on the skull
Current generates a transient magnetic field in the brain inducing electric currents in the cortex that flow parallel to the coil and depolarise neurons
Depending on the frequency, duration of the stimulation, the shape of the coil, and the strength of the magnetic field TMS can activate or suppress activity in cortical regions
Effects of repetitive TMS can outlast the stimulation period for up to 1–2 h.^{24,25}

tDCS (1–2 mA)

Applied through two surface electrodes placed on the skull—eg, for the primary motor cortex (M1), one electrode is placed over M1 and one other over the contralateral supraorbital region
Depending on duration and polarity of stimulation, tDCS enhances or depresses excitability in the stimulated region for minutes to 1–2 h
Does not seem to induce direct neuronal depolarisation like TMS but modulates the activation of sodium and calcium-dependent channels and NMDA-receptor activity, promoting long-term-potential/long-term-depression-like mechanisms

Similarities

Similar duration of effects (up to hours), non-invasive, and under presently used parameters and guidelines safe

Differences

Possible mechanisms underlying their effects; TMS equipment is more expensive, allows more focal stimulation, protocols are better established; TMS applied with millisecond-level accuracy whereas tDCS is applied over minutes
tDCS is easier to use in double-blind or sham-controlled studies, and more easily applied simultaneous to cognitive and motor training protocols

Safety

rTMS can cause seizures, but strict rules of use and training protocols have made them rare³⁶
Patients with a history of seizures should be excluded from studies using this technique
Safety studies are needed for tDCS; brief tingling sensations are common, rarely accompanied by redness under the electrodes^{27,35,38}
Transient headaches have been described for both interventions

Remote effects

Application of both of these techniques to one cortical region will probably influence distant cortical or subcortical sites through trans-synaptic effects
In patients with brain lesions, expected models of current flow elicited by both of these techniques may differ from those in healthy brains³⁷

Non-specific effects on attention, fatigue, discomfort, or mood

Specificity of both of these techniques on cognitive or motor measures may be influenced by possible concomitant effects on attention, fatigue, discomfort, or mood

Study design and experimental hypotheses

Given these characteristics, similarities, and differences the choice of technique to modulate cortical function is highly dependent on study design and experimental hypotheses

task performance³³ and improved acceleration of force production³⁴ compared with sham in the absence of effects on maximum force and finger tapping. Studies with cathodal transcranial direct current stimulation, a




Figure: Targets for intervention strategies based on possible pathophysiological mechanisms
 Movements of the paretic hand are associated with unbalanced interhemispheric inhibition targeting the motor cortex of the affected hemisphere in patients with subcortical stroke.¹ Two interventional approaches might normalise this pattern leading to improvements in motor function: upregulation of excitability of the motor cortex of the affected hemisphere^{29–31} and downregulation of excitability of the motor cortex in the intact hemisphere.^{31,33,34}

form of stimulation that downregulates excitability in the stimulated cerebral cortex, showed results comparable to those achieved with repetitive transcranial magnetic stimulation.³¹ These studies reported improvements in tasks that mimic activities of daily living with the paretic hand, which lasted for up to 25 min. One theoretical advantage of this approach over stimulation of the affected hemisphere is its application to healthy neural structures. Direct comparison of the effects of both strategies is inconclusive.

Different mechanisms not mutually exclusive could contribute to the behavioural gains elicited by non-invasive cortical stimulation. Upregulation of excitability in $M1_{\text{lesioned hemisphere}}$ and downregulation in $M1_{\text{intact hemisphere}}$ could contribute to correcting abnormalities in interhemispheric inhibition identified after stroke,¹ as proposed in previous reports.^{29–31,33,34} This proposal is consistent with the finding that downregulation of excitability in $M1_{\text{intact hemisphere}}$ may correct abnormally high interhemispheric inhibition from $M1_{\text{intact hemisphere}}$ to $M1_{\text{lesioned hemisphere}}$ to an extent proportional to the magnitude of improvements in the paretic hand.⁴⁷ These findings are in line with interhemispheric models in other domains such as attention, memory, language and somatosensory processing.^{48–52,53} One additional

Search strategy and selection criteria

References for this review were identified by searches of MEDLINE between 1969 and April 2006 and references from relevant articles. The search terms used were “brain stimulation”, “stroke”, “recovery”, “TMS”, “tDCS”, “plasticity”, “imaging”, “interhemispheric inhibition”, “intracortical inhibition”, “interhemispheric competition”. Abstracts and reports from meetings were also included. The final reference list was generated based on originality and relevance to the topics covered in the review.

contributing mechanism, proposed by Khedr and colleagues,³² could be the Hebbian interaction between either form of non-invasive stimulation and rehabilitative training.³²

These results were obtained mostly in patients with chronic stroke with moderate impairment (group size between six and 20 patients), as the primary outcome measures for these proof of principle studies were relatively complex skilled movements.^{29–31,33,34} However, Khedr and colleagues³² assessed the effects of repetitive transcranial magnetic stimulation in moderately impaired subacute patients (52 patients) starting with the treatment 1–2 weeks after the stroke. It is important to explore in more detail the usefulness of these methods in patients with poor motor recovery, for whom there are virtually no therapeutic alternatives.

Where next?

Performance improvements reported so far with transcranial magnetic stimulation and transcranial direct current stimulation have been moderate in magnitude (10–30%) and, depending on the study, relatively transient. Implementation of repetitive stimulation and synchronous application with established rehabilitative treatments may lead to more prominent or longer lasting performance improvements. Areas for future investigation are the optimisation of parameters and sham controls, the role of lesion site, time from stroke, drugs and impairment levels on response magnitude, and duration. The successful implementation of these techniques as interventional strategies for these different patient groups will rely on improved understanding of underlying neuronal correlates of functional recovery.^{7,54} Furthermore, from a mechanistic viewpoint, more attention should be paid to possible influences of transcranial magnetic stimulation and transcranial direct current stimulation on subcortical and spinal networks. Additionally, it will be important to determine the extent to which these stimulation techniques influence different types of motor tasks commonly used in neurorehabilitation trials. Despite these limitations, proof of principle studies indicate that non-invasive methods represent powerful approaches to modulate brain function in patients with stroke and could possibly contribute to motor recovery in real-life

neurorehabilitative settings. The data discussed above provide preliminary information, crucial for the design of larger scale, double-blind, sham-controlled clinical trials, which are needed to validate this novel experimental approach.

Contributors

Both authors contributed equally to the preparation of this article.

Conflicts of interest

We have no conflicts of interest.

Acknowledgments

FH's research was supported by a grant from the Alexander von Humboldt Foundation (Feodor-Lynen) and by the Intramural Research Program of the National Institute of Neurological Disorders and Stroke, National Institutes of Health.

References

- Murase N, Duque J, Mazzocchio R, Cohen LG. Influence of interhemispheric interactions on motor function in chronic stroke. *Ann Neurol* 2004; **55**: 400–09.
- Kolominsky-Rabas PL, Weber M, Gefeller O, Neundorfer B, Heuschmann PU. Epidemiology of ischemic stroke subtypes according to TOAST criteria: incidence, recurrence, and long-term survival in ischemic stroke subtypes: a population-based study. *Stroke* 2001; **32**: 2735–40.
- Lai SM, Studenski S, Duncan PW, Perera S. Persisting consequences of stroke measured by the Stroke Impact Scale. *Stroke* 2002; **33**: 1840–44.
- Ward NS, Brown MM, Thompson AJ, Frackowiak RS. Neural correlates of motor recovery after stroke: a longitudinal fMRI study. *Brain* 2003; **126**: 2476–96.
- Calautti C, Baron JC. Functional neuroimaging studies of motor recovery after stroke in adults: a review. *Stroke* 2003; **34**: 1553–66.
- Rossini PM, Calautti C, Pauri F, Baron JC. Post-stroke plastic reorganisation in the adult brain. *Lancet Neurol* 2003; **2**: 493–502.
- Ward NS, Cohen LG. Mechanisms underlying recovery of motor function after stroke. *Arch Neurol* 2004; **61**: 1844–48.
- Liepert J, Hamzei F, Weiller C. Motor cortex disinhibition of the unaffected hemisphere after acute stroke. *Muscle Nerve* 2000; **23**: 1761–63.
- Cicinelli P, Pasqualetti P, Zaccagnini M, Traversa R, Oliveri M, Rossini PM. Interhemispheric asymmetries of motor cortex excitability in the postacute stroke stage: a paired-pulse transcranial magnetic stimulation study. *Stroke* 2003; **34**: 2653–58.
- Manganotti P, Patuzzo S, Cortese F, Palermo A, Smania N, Fiaschi A. Motor disinhibition in affected and unaffected hemisphere in the early period of recovery after stroke. *Clin Neurophysiol* 2002; **113**: 936–43.
- Shimizu T, Hosaki A, Hino T, et al. Motor cortical disinhibition in the unaffected hemisphere after unilateral cortical stroke. *Brain* 2002; **125** (pt 8): 1896–907.
- Kinsbourne M. Mechanisms of hemispheric interaction in man. In: Kinsbourne M, Smith WL, eds. *Hemispheric disconnection and cerebral function*. Springfield, IL: Thomas, 1974: 260–85.
- Calford MB, Tweedale R. Interhemispheric transfer of plasticity in the cerebral cortex. *Science* 1990; **249**: 805–07.
- Ferbert A, Priori A, Rothwell JC, Day BL, Colebatch JG, Marsden CD. Interhemispheric inhibition of the human motor cortex. *J Physiol* 1992; **453**: 525–46.
- Sprague JM. Interaction of cortex and superior colliculus in mediation of visually guided behavior in the cat. *Science* 1966; **153**: 1544–47.
- Kapur N. Paradoxical functional facilitation in brain-behaviour research. A critical review. *Brain* 1996; **119** (pt 5): 1775–90.
- Adkins-Muir DL, Jones TA. Cortical electrical stimulation combined with rehabilitative training: enhanced functional recovery and dendritic plasticity following focal cortical ischemia in rats. *Neurol Res* 2003; **25**: 780–88.
- Kleim JA, Bruneau R, VandenBerg P, MacDonald E, Mulrooney R, Pockock D. Motor cortex stimulation enhances motor recovery and reduces peri-infarct dysfunction following ischemic insult. *Neurol Res* 2003; **25**: 789–93.
- Teskey GC, Flynn C, Goertzen CD, Monfils MH, Young NA. Cortical stimulation improves skilled forelimb use following a focal ischemic infarct in the rat. *Neurol Res* 2003; **25**: 794–800.
- Brown JA, Lutsep HL, Weinand M, Cramer SC. Motor cortex stimulation for the enhancement of recovery from stroke: a prospective, multicenter safety study. *Neurosurgery* 2006; **58**: 464–73.
- Pascual-Leone A, Walsh V, Rothwell J. Transcranial magnetic stimulation in cognitive neuroscience—virtual lesion, chronometry, and functional connectivity. *Curr Opin Neurobiol* 2000; **10**: 232–37.
- Paulus W. Transcranial direct current stimulation (tDCS). *Suppl Clin Neurophysiol* 2003; **56**: 249–54.
- Wassermann EM, Grafman J. Recharging cognition with DC brain polarization. *Trends Cogn Sci* 2005; **9**: 503–05.
- Hallett M. Transcranial magnetic stimulation and the human brain. *Nature* 2000; **406**: 147–50.
- Maeda F, Keenan JP, Tormos JM, Topka H, Pascual-Leone A. Interindividual variability of the modulatory effects of repetitive transcranial magnetic stimulation on cortical excitability. *Exp Brain Res* 2000; **133**: 425–30.
- Siebner HR, Rothwell J. Transcranial magnetic stimulation: new insights into representational cortical plasticity. *Exp Brain Res* 2003; **148**: 1–16.
- Gandiga PC, Hummel FC, Cohen LG. Transcranial DC stimulation (tDCS): a tool for double-blind sham-controlled clinical studies in brain stimulation. *Clin Neurophysiol* 2006; **117**: 845–50.
- Nitsche MA, Seeber A, Frommann K, et al. Modulating parameters of excitability during and after transcranial direct current stimulation of the human motor cortex. *J Physiol* 2005; **568** (pt 1): 291–303.
- Hummel F, Celnik P, Giraux P, et al. Effects of non-invasive cortical stimulation on skilled motor function in chronic stroke. *Brain* 2005; **128** (pt 3): 490–99.
- Hummel F, Cohen LG. Improvement of motor function with noninvasive cortical stimulation in a patient with chronic stroke. *Neurorehabil Neural Repair* 2005; **19**: 14–19.
- Fregni F, Boggio PS, Mansur CG, et al. Transcranial direct current stimulation of the unaffected hemisphere in stroke patients. *Neuroreport* 2005; **16**: 1551–55.
- Khedr EM, Ahmed MA, Fathy N, Rothwell JC. Therapeutic trial of repetitive transcranial magnetic stimulation after acute ischemic stroke. *Neurology* 2005; **65**: 466–68.
- Mansur CG, Fregni F, Boggio PS, et al. A sham stimulation-controlled trial of rTMS of the unaffected hemisphere in stroke patients. *Neurology* 2005; **64**: 1802–04.
- Takeuchi N, Chuma T, Matsuo Y, Watanabe I, Ikoma K. Repetitive transcranial magnetic stimulation of contralesional primary motor cortex improves hand function after stroke. *Stroke* 2005; **36**: 2681–86.
- Nitsche MA, Liebetanz D, Antal A, Lang N, Tergau F, Paulus W. Modulation of cortical excitability by weak direct current stimulation—technical, safety and functional aspects. *Suppl Clin Neurophysiol* 2003; **56**: 255–76.
- Wassermann EM. Risk and safety of repetitive transcranial magnetic stimulation: report and suggested guidelines from the International Workshop on the Safety of Repetitive Transcranial Magnetic Stimulation, June 5–7, 1996. *Electroencephalogr Clin Neurophysiol* 1998; **108**: 1–16.
- Wagner T, Fregni F, Eden U, et al. Transcranial magnetic stimulation and stroke: a computer-based human model study. *Neuroimage* 2006; **30**: 857–70.
- Iyer MB, Mattu U, Grafman J, Lomarev M, Sato S, Wassermann EM. Safety and cognitive effect of frontal DC brain polarization in healthy individuals. *Neurology* 2005; **64**: 872–75.
- Johansen-Berg H, Rushworth MF, Bogdanovic MD, Kischka U, Wimalaratna S, Matthews PM. The role of ipsilateral premotor cortex in hand movement after stroke. *Proc Natl Acad Sci USA* 2002; **99**: 14518–23.
- Fridman EA, Hanakawa T, Chung M, Hummel F, Leiguarda RC, Cohen LG. Reorganization of the human ipsilesional premotor cortex after stroke. *Brain* 2004; **127** (pt 4): 747–58.
- Dancause N, Barbay S, Frost SB, et al. Extensive cortical rewiring after brain injury. *J Neurosci* 2005; **25**: 10167–79.

- 42 Jebsen RH, Taylor N, Trieschmann RB, Trotter MJ, Howard LA. An objective and standardized test of hand function. *Arch Phys Med Rehabil* 1969; **50**: 311–19.
- 43 Plewnia C, Lotze M, Gerloff C. Disinhibition of the contralateral motor cortex by low-frequency rTMS. *Neuroreport* 2003; **14**: 609–12.
- 44 Schambra HM, Sawaki L, Cohen LG. Modulation of excitability of human motor cortex (M1) by 1 Hz transcranial magnetic stimulation of the contralateral M1. *Clin Neurophysiol* 2003; **114**: 130–33.
- 45 Kobayashi M, Hutchinson S, Schlaug G, Pascual-Leone A. Ipsilateral motor cortex activation on functional magnetic resonance imaging during unilateral hand movements is related to interhemispheric interactions. *Neuroimage* 2003; **20**: 2259–70.
- 46 Swinnen SP. Intermanual coordination: from behavioural principles to neural-network interactions. *Nat Rev Neurosci* 2002; **3**: 348–59.
- 47 Hummel F, Celnik P, Floel A, et al. Effects of noninvasive cortical stimulation (tDCS) of the intact hemisphere in patients with chronic stroke on (II) interhemispheric inhibition. *Neuroimage* 2005; **26** (suppl 1): S34.
- 48 Floel A, Nagorsen U, Werhahn KJ, et al. Influence of somatosensory input on motor function in patients with chronic stroke. *Ann Neurol* 2004; **56**: 206–12.
- 49 Hilgetag CC, Theoret H, Pascual-Leone A. Enhanced visual spatial attention ipsilateral to rTMS-induced ‘virtual lesions’ of human parietal cortex. *Nat Neurosci* 2001; **4**: 953–57.
- 50 Naeser MA, Martin PI, Nicholas M, et al. Improved picture naming in chronic aphasia after TMS to part of right Broca’s area: an open-protocol study. *Brain Lang* 2005; **93**: 95–105.
- 51 Oliveri M, Rossini PM, Traversa R, et al. Left frontal transcranial magnetic stimulation reduces contralesional extinction in patients with unilateral right brain damage. *Brain* 1999; **122** (pt 9): 1731–39.
- 52 Oliveri M, Bisiach E, Brighina F, et al. rTMS of the unaffected hemisphere transiently reduces contralesional visuospatial hemineglect. *Neurology* 2001; **57**: 1338–40.
- 53 Corbetta M, Kircade MJ, Lewis C, Snyder AZ, Sapir A. Neural basis and recovery of spatial attention deficits in spatial neglect. *Nat Neurosci* 2005; **8**: 1603–10.
- 54 Lotze M, Market J, Sauseng P, et al. The role of multiple contralesional motor areas for complex hand movements after internal capsular lesion. *J Neuroscience* 2006; **26**: 6096–102.